Influenza Surveillance Report

Division of Epidemiology - Disease Surveillance and Investigation

District of Columbia Department of Health

2011-2012 Influenza Season Weeks 18 – 19 (April 29, 2012 – May 12, 2012)

(All data are preliminary and may change as more reports are received)

SUMMARY

- 8 cases of Influenza were reported by hospitals during this reporting period.
- Zero pediatric-deaths associated with Influenza were recorded during this reporting period.
- During this reporting period, one hospitalization for Influenza was recorded, with ten to-date.
- For the 2011-2012 Influenza Season to-date, 92 positive Influenza cases have been reported.

INFLUENZA SURVEILLANCE FROM HOSPITALS & AMBULATORY CARE FACILITIES


District of Columbia hospitals and laboratories report detailed information on cases of Influenza on a daily basis. However, in accordance with CDC guidelines, only Influenza-associated deaths in cases <18 years of age and Novel Influenza A infections are reportable.


The table below summarizes weekly and cumulative cases of Influenza for the 2011-2012 Season. Data are also presented by age group and by number of cases reported weekly. During weeks 18 – 19 (April 29, 2012 – May 12, 2012), there were eight new cases of Influenza reported. To date, the District has received 92 positive Influenza cases among DC residents reported by hospitals.

Surveillance of Influenza Cases Reported By Influenza Type

	Weeks 18 – 19 (29 April 2012 – 12 May 2012)		Cumulative Cases for Weeks 40 – 22 (2 October 2011 – 2 June 2012)	
Influenza A	6	(75%)	60	(65.2%)
Influenza B	2	(25%)	25	(27.2%)
Influenza A/B	0	(0%)	3	(3.3%)
Influenza (not typed)	0	(0%)	4	(4.3%)
Total	8*	(100%)	92*	(100.0%)

^{*}Includes results from Rapid Diagnostic Testing, Viral Culture, RT-PCR, Serology, and Immunofluorescence.





RAPID DIAGNOSTIC TESTING

Rapid Diagnostic Tests are screening tests used to detect the Influenza virus in a short period of time. While initially less accurate than PCR and viral culture, rapid diagnostics are more accurate as the Influenza season progresses. During weeks 18 – 19, 107 tests out of a total of 112 were performed using Rapid Diagnostic Testing in clinical laboratories. Of these, eight (7.4%) positive Influenza specimens were identified during weeks 18 – 19 using rapid diagnostics. The other five tests were performed using viral culture.

Weeks: 18 – 19 (29 April 2012 – 12 May 2012)				
No. of specimens tested	107			
No. of positive specimens (%)	8 (7.4%)			
Positive specimens by type/subtype				
Influenza A	6 (75%)			
Influenza B	2 (25%)			
Influenza A/B	0 (0%)			
Influenza – unknown type	0 (0%)			

INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE

Sentinel surveillance for ILI consists of three outpatient reporting sites for the District of Columbia. The sentinel surveillance sites report the total number of ILI cases encountered per week and the total number of patients seen at the clinic during that same week. For this system, ILI is defined as the existence of fever (temperature of 100°F [37.8°C] or greater) and a cough and/or a sore throat in the absence of a known cause other than Influenza.

For weeks 18 – 19, sentinel providers reported that 101 out of 3,738 visits (2.7%) met the criteria for ILI.

Sentinel Surveillance ILI Activity for Washington, DC


Week of	Activity *			
April 29 – May 5	Sporadic			
May 5 – May 12	Sporadic			

*No Activity – overall clinical activity remains low and there are no lab confirmed Influenza cases;

Sporadic – isolated lab confirmed Influenza cases reported and ILI activity is not increased:

Local – increased ILI activity and recent lab confirmed Influenza cases. As the District of Columbia is not a state, this is the highest level of ILI activity it can report.


Influenza-Like Illness Reported by MMWR Weekly October 2, 2011 - June 2, 2012



SYNDROMIC SURVEILLANCE FOR THE DISTRICT OF COLUMBIA


The Electronic Surveillance System for Early Notification of Community Epidemics (ESSENCE) is a valuable tool for early detection of naturally occurring and induced health events throughout the National Capital Region (NCR). Syndromic surveillance involves the parsing of a patient's emergency department (ED) chief complaints into predetermined syndromes and subsyndromes. Influenza-like illness (ILI) is a subsyndrome in ESSENCE that is defined by the presence of "Influenza," or "Fever" plus "Cough" or "Sore Throat." By tracking ILI, the severity of the Influenza season can be monitored and compared to that of previous years.

The percentage of visits to EDs by week for ILI is shown in orange for the current 2011-2012 Influenza Season. The percentage of ILI visits made to hospital EDs for the past five Influenza seasons plotted for are also comparison. Data since weeks 10-11 are not available in ESSENCE, and therefore the graph depicts only data through week 9. The graph will be updated when data are available.




- * CDC defines the Influenza season as beginning in October and ending mid-May (Weeks 40-21)
- ** The 2009-2010 Influenza Season featured an unusual pattern of illness with the emergence of Novel H1N1 Influenza A.

The percentage of visits to EDs by month for ILI is shown in orange for the current 2011-2012 Influenza Season, plotted against the past five Influenza Seasons. The data for March and April will be reported when data are available.




** The 2009-2010 Influenza Season featured an unusual pattern of illness with the emergence of Novel H1N1 Influenza A.

The percentage of visits to EDs for ILI is shown in blue for the current 2011-2012 Influenza Season by MMWR week. The mean percentage of ILI visits made to hospital EDs for the past five Influenza seasons is also plotted for comparison. Data since weeks 10-11 are not available in ESSENCE, and therefore the graph depicts only data through week 9. The graph will be updated when data are available.



** The 2009-2010 Influenza Season, included in the 5-Season mean, featured an unusual pattern of illness with the emergence of Novel H1N1 Influenza A.

The percentage of visits to EDs for ILI is shown in blue for the current 2011-2012 Influenza Season by month, plotted against the mean from the past five Influenza Seasons. The data for March and April will be reported when data are available.



^{**} The 2009-2010 Influenza Season, included in the 5-Season mean, featured an unusual pattern of illness with the emergence of Novel H1N1 Influenza A.

INFLUENZA TESTING BY THE DISTRICT OF COLUMBIA PUBLIC HEALTH LABORATORY (DC PHL)

The DC PHL subtypes human isolates to monitor the circulating strains of Influenza. The isolates are submitted to the DC PHL by hospitals and commercial laboratories. To date, twenty six out of fifty-eight specimens sent to the DC PHL have tested positive for Influenza. Cumulatively, fifteen of these isolates were subtyped as Influenza A/H3, six were subtyped as Influenza A/H1, and five were Influenza B.

DC PHL Surveillance of Influenza Cases Reported By Influenza Subtype


	Weeks 18 – 19* (29 April 2012 – 12 May 2012)	Cumulative Cases for Weeks 40 – 22 (2 Oct 2011 – 2 June 2012)	
Number of specimens	6	58	
Number of specimens positive for Influenza:	3/6 (50%)	26/58 (44.8%)	
• Influenza A	2/3 (66.7%)	21/26 (80.8%)	
■ H1	0/2 (0%)	6/21 (28.6%)	
■ нз	2/2 (100%)	15/21 (71.4%)	
Influenza B	1/3 (33.3%)	5/26 (19.2%)	

^{*} Week designation for this data is based on the date that the isolate was received by DC PHL.

NATIONAL INFLUENZA ASSESSMENT

The CDC's weekly seasonal Influenza surveillance report for weeks 18 - 19 noted that Influenza-like illness (ILI) activity among outpatients in the United States remained elevated but on the decline nationally and in most regions. The proportion of deaths due to Influenza and pneumonia in the US was below the epidemic threshold. Two Influenza-associated pediatric deaths were recorded in the US during week 18, one due to untyped Influenza A and one due to Influenza B. Two pediatric deaths occurred during week 19, one due to untyped Influenza A and one due to Influenza B. Of the 579 respiratory specimens that tested positive during weeks 18-19, 58.7% were Influenza A viruses. Of the Influenza A samples, 51.8% were Influenza A (H3), 10.9% were Influenza A (2009 H1N1), and 37.9% of the viruses were not subtyped.

Influenza Positive Tests Reported to CDC by U.S. WHO/NREVSS Collaborating Laboratories, National Summary, 2011-12



Get Vaccinated!

To find an Influenza vaccine provider, visit the District of Columbia Flu Resource Center at http://doh.dc.gov/doh/cwp/view,a,1370,q,604320.asp



For additional information about Influenza and Influenza activity in the United States, please visit: http://www.cdc.gov/flu/index.htm.

Questions about Influenza in the District of Columbia or this report should be directed to the Division of Epidemiology - Disease Surveillance and Investigation at (202) 442-8141 or email gabrielle.ray@dc.gov.